

TRADITIONAL FERMENTED FISH PRODUCTS IN INDONESIA

by

HARI EKO IRIANTO

Slipi Research Station for Marine Fisheries
Jalan Petamburan VI, Jakarta 10260, INDONESIA

and

GIYATMI IRIANTO

Food Technology Department, Sahid University
Jalan Prof. Supomo No.84, Jakarta 12870, INDONESIA

ABSTRACT

Fermented fish products have an important role in stimulating appetite by providing unique aromas and flavours. Many kinds of fermented fish products can be found in various parts of Indonesia. Raw materials include whole fish, comminuted fish and viscera from marine and fresh water fishes. The products are peda, jambal roti, kecap ikan, terasi, ikan tukai, bekasang, bekasam, naniura, picungan, and cincaluk.

INTRODUCTION

Traditional processing makes a significant contribution to fish preservation in Indonesia as about 50 percent of the total marine catch is processed and traditional methods account for more than 40 percent of that.

One of the traditional fish processing methods is fermentation and many kinds of fermented fish products can be found in parts of Indonesia. Those products are able to provide specific unique characteristics. In fermented processing, there is a transformation of organic substances into simpler compounds either by the action of micro-organisms or by the action of enzymes from the fish (Beddows, 1985). At the same time, the fish odour of the starting material is changed to the flavour and aromas of certain foods (Bukholder *et al.*, 1968). Enzymes are most significant in changing the texture and producing some of the flavour but micro-organisms aid in the development of aroma and flavour (Beddows, 1985).

Consumption of fermented fish products in Indonesia is mainly to introduce specific flavours which encourage the appetite. The many fermented fish products available in Indonesia provide many different flavours which meet various consumer appetites. Unfortunately, most of the fermented products are only available locally. Thus, a fermented product in one province will be very difficult to find in other provinces.

This paper mainly addresses fermented fish products found in Indonesia. Due to lack of scientific information about most of the products, the paper emphasizes processing technologies, which are traditionally employed by processors.

PRODUCT TYPES

Based on the information which has been compiled, Indonesian fermented fish can be divided in several ways, such as in terms of raw materials, fermentation types and final product forms.

a. Raw Materials

1. fresh water fish: bekasam
2. marine fish:
 - 2.1. whole gutted fish: peda, jambal roti, ikan tukai, cincaluk, picungan, and naniura
 - 2.2. pounded fish/shrimp: terasi
 - 2.3. visceral: bekasang

b. Types of Fermentation

1. fish and salt fermentation: peda, jambal roti, ikan tukai, bekasang, terasi, and kecap ikan
2. fish, carbohydrate and salt fermentation: bekasam, cincaluk, naniura, and picungan

c. Final Product Forms

1. dried fish: peda, jambal roti, and ikan tukai
2. moist fish: bekasam, naniura, and picungan
3. lumped pounded fish/shrimp: terasi
4. liquid/semi-liquid: kecap ikan, bekasang, and cincaluk.

FERMENTED FISH PRODUCTS

Peda

Peda is mainly found in Java. This product is processed using mackerel (*Rastrelliger negletus*) as raw material. Fresh water fish processed into peda did not result in products with similar quality (Sukarsa, 1979). Nutritional quality of peda can be seen in Table 1.

Table 1. Nutritional quality of Peda.

| Parameter | Amount in 100 g sample |
|----------------------|------------------------|
| Moisture content (g) | 46 |
| Energy (Calorie) | 156 |
| Protein content (g) | 28 |
| Fat content (g) | 4 |
| Ca (mg) | 174 |
| P (mg) | 316 |
| Fe (mg) | 3.1 |
| Vitamin A (IU) | 110 |
| Vitamin B1 (mg) | trace |
| Vitamin C (mg) | trace |

Source: Soedarmo and Sediaoetama (1977)

The basic method of peda processing is a salting process with two steps. The first salting step takes several weeks to develop its characteristic flavour and texture, and this is followed by a maturation phase. Salting is carried out using a 1:3 salt to fish ratio. Fish and salt are arranged in layers alternately in a salting tank. By the end of the process, the fish are soaked in saturated brine pickle with coarse salt remaining at the bottom and the top covering the fish. This salting process normally takes 3 days. Maturation is performed by mixing the fish and salt thoroughly. The amount of salt used is a one third of fish weight. During maturation, the fish are kept in bamboo baskets, the process taking around 1-2 weeks (Rahayu, 1992). Better products can be obtained by using eviscerated fish (Hanafiah, 1987) and without salt addition in the second fermentation (Hanafiah, 1987; Irianto, 1990; Irianto and Brooks, 1994).

Investigations conducted by Nur and Sjahri (1979) indicate that fish viscera and vacuum encourage the fermentation process. Use of a starter made from matured peda has a similar effect to fish viscera. Sjachri and Nur (1979) improved pedah quality by introducing antifungals and antioxidants during processing.

Microbiological studies on peda conducted by Hanafiah (1987) revealed Gram-positive cocci predominated and some were identified as lactic acid bacteria. Isolation of halo-tolerant bacteria from peda by Suwandi (1988) concluded that bacterial growth in peda was characterized as Gram positive cocci, which were non-motile, aerobes or facultative aerobes, catalase positive, non-indole producers and oxidase negative. They can utilize citrate as the only carbon source, ferment glucose and show proteolytic activity. Some of them are able to reduce nitrate. These bacteria can be classified as mesophiles and require a pH of 6-8. They show variations in salt tolerance and can be divided on the basis of salt tolerance as weak, moderate and halotolerant bacteria. Idawati (1996) showed that lactic acid bacteria isolated from peda were homo-fermentative and hetero-fermentative *Lactobacillus* sp., *Leuconostoc* sp., and *Streptococcus* sp.

Jambal Roti

Jambal roti is processed from marine catfish (*Arius thalassinus*). Production centres in Java are Pekalongan, Cilacap, Cirebon and Pangandaran. Basically, processing methods applied in all production places are the same, but each location has their own specific ways to process jambal roti.

In the processing method employed by processors in Cirebon, the fish are first beheaded and eviscerated. They are then washed and soaked in fresh water for 24 hours. After draining, the fish are arranged in a basin. Each layer of fish is sprinkled with salt and the bellies are also filled with salt. Total amount of salt used for salting is 30-35% of fish weigh and the salting process takes around 24 hours. After salting, the fish are freed from excessive salt and washed. Clean fish are soaked in fresh water for 15-20 minutes and subsequently split into butterfly form. The split fish is sun dried for 3-5 days (Burhanuddin *et al.*, 1987).

Fish which has been iced usually produces a worse quality product than fish without icing. Nuraniekmah (1996) investigated a new method for processing jambal roti from iced fish by soaking the fish in warmer water to bring the temperature back to normal before processing. The higher the water temperature, the shorter the soaking period will be. Soaking the fish at 40°C needs 60 minutes to reach normal temperature.

Erwan (1992) developed a modified method of jambal roti processing by soaking the fish in 30% coconut sugar solution before salting. He also found that 20% of fish weight was enough for salting. Proximate composition, salt content and pH of jambal roti noted by Nuraniekmah (1996) can be seen in Table 2.

Table 2. Proximate composition, salt content and pH of jambal roti.

| Parameter | |
|----------------------|---------------|
| Moisture content (%) | 49.27 - 49.68 |
| Protein content (%) | 54.17 - 61.86 |
| Fat content (%) | 0.69 - 1.19 |
| Ash content (%) | 34.93 - 38.80 |
| Salt content (%) | 7.38 - 8.53 |
| pH | 6.57 - 6.91 |

Source: Nuraniekmah (1996)

Terasi

Terasi is consumed in small quantities as a flavour. The product is not only for local consumption, but is also exported, mainly to the Netherlands and Suriname, in powder form. Terasi is usually made from planktonic shrimp "rebon" (*Atyas* sp. or *Mycis* sp.) (Budhyatni *et al.*, 1982). Terasi processors can be found in Java and Sumatera islands. According to Yunizal (1998) terasi can be processed in two ways, i.e. (1) with salt only and (2) with salt and other ingredients.

In the processing of terasi with salt only, firstly rebon is washed, drained and dried, until half dried. During drying, impurities, such as small fish, mussel shells and coral, are removed. After that, semi dried rebon is sifted to separate sand and other undesirable materials. The rebon is then left overnight at ambient temperature and pounded the next day. During the first pounding, salt is added (around half of the total salt required during processing). Total amount of salt used in terasi processing is 2-5% of rebon weight which should be added as solution. Pounded rebon is sun dried and subsequently kept in a container at ambient temperature for 2-3 days. The stored rebon is then pounded for a second time, while the remaining salt is added. After that, the pounded rebon is sun dried and kept at ambient temperature for 2-3 days until soft. It is then ground by many passes through a meat grinder until fine. Fine rebon is formed in cubes or cylinders of 1 kg weight and subsequently fermented for a week or more at ambient temperature (Yunizal, 1998). Proximate composition and important mineral contents of this type of terasi can be seen in Table 3.

Table 3. Proximate composition, energy, and mineral contents of terasi.

| Parameter | Amount in 100 g sample |
|----------------------|------------------------|
| Moisture content (g) | 40 |
| Energy (Calories) | 174 |
| Protein content (g) | 30 |
| Fat content (g) | 3.5 |
| Ca (mg) | 100 |
| P (mg) | 250 |
| Fe (mg) | 3.1 |

Source: Soedarmo and Sediaoetama (1977)

Processing method of terasi with salt and other ingredients is similar to the processing method of terasi with salt only. Specifically, in the second pounding, the salt solution is mixed with coconut sugar and tamarind prior to be added to rebon. For 10 kg fresh rebon, 200g and 250g respectively of tamarind and coconut sugar are used. The additional ingredients accelerate the fermentation process (Yunizal, 1998).

Budhyatni *et al.* (1982) investigated micro-organisms in terasi powder during ambient storage. Results showed that the number of *Lactobacillus* sp was constant during storage. Pathogenic bacteria: *Staphylococcus* sp., *Bacillus* sp., and *Proteus* sp. were present, but *Salmonella* sp., *Clostridium* sp., *Vibrio* sp. and *E. coli* were absent.

Kecap Ikan

Kecap ikan (fish sauce) is not really popular in Indonesia due to strong competition from soy sauce, but is well recognized in West Kalimantan Province (Mulyokusumo, 1974). In East Java, kecap ikan is processed from oil sardine (*Sardinella lemuru*) (Putro, 1993).

Kecap ikan is traditionally produced by fermentation using high salt concentrations. Generally fermentation takes a long time to complete (Suparno and Silowati, 1982).

In the processing of kecap ikan described by Putro (1993), fish are firstly washed and minced. Minced fish are mixed with salt (25-30% of fish weight) and allowed to ferment for 10-12 months. The fermented mixture is then filtered and finally brown sugar and spices are added. Results of chemical analysis of commercial kecap ikan can be seen in Table 4.

Table 4. Chemical composition of kecap ikan.

| Parameter | Kecap Ikan A | Kecap Ikan B |
|--------------------------|--------------|--------------|
| Moisture content (%) | 66.67 | 76.89 |
| Ash content (%) | 23.5 | 21.95 |
| Salt content (%) | 21.16 | 11.60 |
| Protein content (%) | 10.17 | 10.51 |
| Fat content (%) | 0.50 | 0.70 |
| Carbohydrate content (%) | 1.50 | 0.30 |
| Energy (calories/g) | 5.41 | 5.24 |

Source: Poernomo et al. (1984)

Idawati (1996) produced kecap ikan from the brine waste of salting peda. The brine waste was fermented for a month. Identification of lactic acid bacteria isolated from that kecap ikan revealed that bacteria involved in the fermentation of kecap ikan were homo fermentative *Lactobacillus* sp, *Pediococcus* sp and *Streptococcus* sp.

Kumalarningsih (1986) studied the improvement of fish sauce quality by introducing bromeline and papain to prepare fish hydrolysate as well as the use of *Brevibacterium linens* and *Micrococcus* sp. According to the formation of amino-N, the incorporation of *Brevibacterium linens* strain C and *Micrococcus* were considered to be promising.

Ikan Tukai

Ikan tukai is a traditional fermented fish product which is only found in West Sumatera. Ikan tukai is also called *lauak tukai* or *ikan sambal lado*, and is mostly processed from barracuda (*Sphyranea* sp.).

The traditional processing method of ikan tukai is unique. Barracuda is washed and soaked in 20% brine for around two hours. After draining, the fish are dried for a day. Dried fish are then wrapped with taro leaves. The wrapped fish are kept underground for two days to allow fermentation and then sun dried until dry. Chemical characteristics of commercial ikan tukai can be seen in Table 5.

Table 5. Results of chemical analyses of ikan tukai.

| Analyses | |
|----------------------|--------|
| Moisture content (%) | 51.01 |
| Salt content (%) | 5.05 |
| pH | 6.93 |
| TVN (mg/100g) | 113.22 |
| VRS (meq/g) | 40.97 |

Source: Effendi (1995)

Effendi (1995) processed ikan tukai using a modified method, in which fermentation and drying were carried out in vacuum incubator and oven respectively. His microbiological studies concluded that bacteria

having a significant role in the fermentation process of ikan tukai were *Micrococcus* sp., *Pediococcus* sp., *Lactobacillus* sp, *Pseudomonas* sp and *Staphylococcus* sp.

Bekasang

Bekasang is a traditional product in North Sulawesi and the Moluccas and is processed from the viscera of skipjack (*Katsuwonus pelamis*), which is waste from *cakalang fufu* (smoked skipjack) processing in North Sulawesi and *cakalang asar* (smoked skipjack) processing in the Moluccas.

Processing methods of bekasang employed by processors in Manado-North Sulawesi are as follows. Skipjack viscera obtained from cakalang fufu processors is washed and mixed with salt at a ratio of 2.5:1. The mixture of viscera and salt is kept for a week to allow fermentation. After fermentation ceases, the fermented viscera are boiled for 2 hours and filtered using gauze. The filtrate is bottled and ready to sell (Wudianto *et al.*, 1996).

Subroto *et al.* (1984) introduced a mincing treatment to viscera before mixing with salt, this resulted in better quality bekasang. Setiabudi *et al.* (1985) investigated salt levels in bekasang processing by varying salt addition from 10 to 20%. Conclusions from the results were that the higher salt addition, the better bekasang quality obtained. Comparison of protein content and pH of experimental bekasang with commercial bekasang can be seen Table 6.

Table 6. Protein content and pH of experimental and commercial bekasang.

| Samples of Bekasang | Protein content (%) | pH |
|------------------------|---------------------|------|
| Experimental bekasang: | | |
| - from minced viscera | 37.95 | 5.65 |
| - from whole viscera | 36.52 | 5.72 |
| Commercial bekasang | 22.37 | 5.77 |

Source: Subroto *et al.* (1984)

Bekasam

Bekasam is a fermented fish product processed from fresh water fish which can be found in South Sumatera and Central Kalimantan. Bekasam is served by mixing with chilli and sugar. In bekasam processing, inspite of the salt addition, carbohydrate sources are also incorporated to stimulate the growth of lactic acid bacteria by decomposing into simpler compounds. Carbohydrate sources used are cooked rice, roasted rice and sticky rice Murtini (1992).

Processing of bekasam from common carp (*Cyprinus carpio* Linn) was described by Murtini (1992). Firstly, the fish are beheaded, descaled and eviscerated. They are then cut into butterfly forms and washed. Washed fish are eventually soaked in 16% brine solution for 48 hours prevented from floating by weights. The fish are then drained and cooked rice and sticky rice, up to 50% and 25% of fish weight respectively, are added. Finally the mixture of fish and rice is sealed in plastic jars and allowed to ferment for a week or more. Proximate composition and salt content of common carp bekasam are shown in Table 7.

Table 7. Proximate composition and salt content of common carp bekasam.

| Parameter | |
|----------------------|-------|
| Moisture content (%) | 66.95 |
| Ash content (%) | 5.76 |
| Protein content (%) | 4.81 |
| Fat content (%) | 14.95 |
| Salt content (%) | 5.72 |

Source: Murtini (1992)

In order to improve bekasam quality, Murtini *et al.* (1997) added the liquids of cabbage and Chinese leaf pickles as lactic acid bacteria sources into spotted gouramy bekasam. Sensory evaluation indicated that the best bekasam was produced by the addition of Chinese leaf pickle. All bekasam in this study was organoleptically still acceptable after 8-weeks storage.

A similar product called *Naniura* is found in Riau and North Sumatera provinces. In *naniura* processing, the fish are firstly soaked in lemon juice or 25% acetic acid solution for three hours. After that, ground boiled rice is added and the fish is then packed and allowed to ferment.

Another similar product to bekasam is *Picungan* which can be found in West Java. This product is processed using marine fish. *Picungan* seeds are used as a carbohydrate source. The seeds are cut into small pieces before mixing with fish and salt.

Cincaluk

Cincaluk is a traditional fermented fish product from Riau Province. This product is usually processed from *rebon*.

There is no prescribed method to process cincaluk. In a method employed by processors in Bengkalis, fresh shrimp are mixed with boiled rice and salt in a pan. For 1 kg shrimp, the rice added is around 200-300g, while salt is approximately 300g. The pan is then sealed with the lid to avoid air entry and kept for 4 days until liquid release. After that the mixture is put into bottles and eventually sealed tightly. Proximate composition and other chemical properties of cincaluk processed using this method can be seen in Table 8.

Another method to process cincaluk is by mixing shrimp with tapioca flour, salt and sugar at the ratio of 20:1:1:1. In its processing, shrimp are descaled and then washed. Tapioca flour is dissolved in water, gelatinized and allowed to cool. Shrimp are then mixed thoroughly with salt, sugar and gelatinized tapioca flour. The mixture is filled to washed bottles and sealed firmly and fermented at ambient temperature for 1-2 weeks.

Table 8. Proximate composition and other chemical properties of cincaluk.

| Parameter | |
|-------------------------|-------|
| Moisture content (%) | 69.76 |
| Ash content (%) | 12.43 |
| Protein content (%) | 16.23 |
| Fat content (%) | 1.57 |
| Salt content (%) | 10.11 |
| Lactic acid content (%) | 2.34 |
| pH | 4.82 |

CONCLUSIONS

Indonesian fermented fish products vary in form, raw material and fermentation type. Most of them have not been studied in detail, thus scientific information relating to those products is difficult to find. More studies identify lactic acid bacteria involved in the fermentation which suggests improved product quality can be achieved by using selected lactic acid bacteria.

REFERENCES

- Beddows, C.G.. 1985. Fermented fish and fish products. *In* Microbiology of fermented foods vol 2. Edited by Wood, B.J.B., Elsevier Applied Science. London. p.1-39.
- Budhyatni, S. , Murtini, J.T. and Peranginangin, R. 1982. The microflora of terasi powder. Laporan Penelitian Teknologi Perikanan 16: 25-33 (*In Indonesian*).
- Burkholder, L. , Burkholder, P.R. , Chu, A. , Kostyk, N. and Roels, O.A. 1968. Fermentation. Food Tech 22: 1278-84.
- Burhanuddin, S. , Martosewojo, S. Djamali, A. and Hutomo, M. 1987. Resources of marine catfish in Indonesia. Pusat Penelitian dan Pengembangan Oseanologi Nasional - LIPI. Jakarta (*In Indonesian*).
- Efendi, Y. 1995. Preliminary study on the processing of ikan tukai. *In* Proceeding of the First Symposium of Indonesia Fisheries. Jakarta, August 25-27, 1993. Pusat Penelitian dan Pengembangan Perikanan. Jakarta p. 152-63.
- Erwan, M. 1992. Effects of sugar and salt concentration on jambal roti quality. Faculty of Fisheries-Bogor Agricultural University. Bogor (*In Indonesian*).
- Hanafiah, T.A.R. 1987. Factors affecting quality of pedah siam. Master Thesis. University of Washington, Seattle.
- Idawati. 1996. Isolation and selection of lactic acid bacteria having antibacterial activities from peda and kecap ikan. Sarjana Thesis. Faculty of Agricultural Technology-Bogor Agricultural University. Bogor (*In Indonesian*).
- Irianto, H.E. 1990. Studies on the processing of pedah, a traditional Indonesian fermented fish product. Diploma Thesis. Massey University. New Zealand.
- Irianto, H.E. and Brooks, J.D. 1994. Investigation on the optimum conditions in pedah processing. Jurnal Penelitian Pasca Panen Perikanan 81: 18-29.
- Kumalaningsih, S. 1986. Incorporation of proteolytic enzymes and bacteria in the fermentation of kecap ikan of oil sardine (*Sardinella* sp.). Ph.D Dissertation. Brawijaya University. Malang (*In Indonesian*).
- Mulyokusumo. 1974. Soy sauce, peanut sauce, fish sauce. Terate. Bandung (*In Indonesian*).
- Murtini, J.T. 1992. Common carp bekasam. *In* Compilation of research results of fishery post-harvest. Edited by Suparno, Nasran, S. and Setiabudi, E. Pusat Penelitian dan Pengembangan Perikanan. Jakarta p.135-6 (*In Indonesian*).

- Murtini, J.T , Yuliana, E. , Nurjanah and Nasran, S. 1997. Effects of addition of lactic acid bacteria starter in the processing of spotted gouramy (*Trichogaster trichopterus*) bekasam on its quality and shelflife. *Jurnal Penelitian Perikanan Indonesia* III (2): 71-82 (*In Indonesian*).
- Nur, M.A. and Sjachri, M. 1979. Processing of traditional product. I. Effects of several treatments on physical and chemical properties of peda processed in laboratory. *In Laporan Lokakarya Teknologi Pengolahan Ikan Secara Tradisional*. Jakarta, 26 Pebruari - 1 Maret 1979. Lembaga Penelitian Teknologi Perikanan. Jakarta. p.91-4 (*In Indonesian*).
- Nuraniemah, S.R. 1996. Effects of soaking temperature on proteolytic enzyme activity and bacteria growth of jambal roti made of marine catfish (*Arius thalassinus*). Sarjana Thesis. Faculty of Fisheries-Bogor Agricultural University. Bogor (*In Indonesian*).
- Poernomo, A. , Suryaningrum, T.D. Ariyani, F. and Putro, S. 1984. Studies on the nutritive value and microbiology of traditional fishery products. *Laporan Penelitian Teknologi Perikanan* 30: 9-19 (*In Indonesian*).
- Putro, S. 1993. Fish fermentation technology in Indonesia. *In Fish fermentation Technology*. Edited by Lee, C.H., Steinkraus, K.H. and Reilly, P.J.A. United Nations University Press. Tokyo p. 107-28.
- Rahayu, S. 1992. Processing of peda. *In Compilation of research results of fishery post-harvest*. Edited by Suparno, Nasran, S. and Setiabudi, E. Pusat Penelitian dan Pengembangan Perikanan. Jakarta p.133-4 (*In Indonesian*).
- Setiabudi, E. , Subroto, W. and Bustaman, S. 1985. Effects of salt content on the quality of bekasang during fermentation process. *Laporan Penelitian Teknologi Perikanan* 46: 11-5 (*In Indonesian*).
- Soedarmo, P. and Sediaoetama, A.D. 1977. Nutrition science. Penerbit Dian Rakyat (*In Indonesian*).
- Subroto, W. , Setiabudi, E. and Bustaman, S. 1984. Preliminary study on the production of bekasang. *Laporan Penelitian Teknologi Perikanan* 26: 9-16 (*In Indonesian*).
- Sukarsa, D.R. 1979. Processing of peda from fresh water fish. *In Laporan Lokakarya Teknologi Pengolahan Ikan Secara Tradisional*. Jakarta, 26 Pebruari - 1 Maret 1979. Lembaga Penelitian Teknologi Perikanan. Jakarta. p.94-100 (*In Indonesian*).
- Suparno and Silowati, T. 1982. Preparation of fish sauce from mackerel (*Rastrelliger* spp) by acid hydrolysis. *Laporan Penelitian Teknologi Perikanan* 20: 29-36 (*In Indonesian*).
- Suwandi, I. 1988. Studies on the physiological characteristics of halotolerant bacteria isolated from peda. Sarjana Thesis. Bogor Agricultural University. Bogor (*In Indonesian*).
- Syachri, M. and Nur, M.A. 1979. Processing of traditional product. II. Effects of the use of anti fungal (sorbic acid) and anti oxidant (BHA) on chemical characteristics of peda. *In Laporan Lokakarya Teknologi Pengolahan Ikan Secara Tradisional*. Jakarta, 26 Pebruari - 1 Maret 1979. Lembaga Penelitian Teknologi Perikanan. Jakarta. p.162-166 (*In Indonesian*).
- Wudianto, Naamin, N., Susanto, K., Irianto, H.E. and Pranowo, S. A. 1996. A Fishery and socio-economic survey in MCMA of Karakelong-Manado, North Sulawesi Pusat Penelitian dan Pengembangan Perikanan. Jakarta (*In Indonesian*).
- Yunizal. 1998. Processing of shrimp terasi. *Warta Penelitian dan Pengembangan Pertanian* XX (1): 4-6 (*In Indonesian*).